



Nadkarni, A; Bhat, B; Ebrahim, S; Patel, V (2013) The course and outcome of alcohol use disorders in men in Goa: A population- based follow-up study. Indian journal of psychiatry, 55 (4). pp. 376-379. ISSN 0019-5545

Downloaded from: <http://researchonline.lshtm.ac.uk/1410530/>

DOI:

#### Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>



# Indian Journal of Psychiatry

OFFICIAL PUBLICATION OF THE INDIAN PSYCHIATRIC SOCIETY

ISSN 0019-5545

Volume 55, Number 4

October-December 2013

Now indexed  
with PubMed

## A TRIBUTE

Prof. J. K. Trivedi  
ROY ABRAHAM KALLIVAYALIL, PRONOB KUMAR DALAL

311

## EDITORIAL

Psychosomatic paradigms in psoriasis – Psoriasis, stress and mental health

T. S. SATHYANARAYANA RAO, K. H. BASAVARAJ, KEYA DAS

313

## GUEST EDITORIALS

Sexual abuse in women with special reference to children: Barriers, boundaries and beyond  
P. B. BEHERE, T. S. SATHYANARAYANA RAO, AKSHATA N. MULMULE

Loneliness – A disease?

SARVADA CHANDRA TIWARI

## REVIEW ARTICLE

Group psychotherapies for depression in persons with HIV – A systematic review

ABHIJIT RAMANNA HONAGODU, MURALI KRISHNA, RAJESH SUNDARACHAR, PETER LEPPING

## ORIGINAL ARTICLES

Assessment of cognition in non-affected full biological siblings of patients with schizophrenia  
ROHIT GARG, J. K. TRIVEDI, P. K. DALAL, ANIL NISCHAL, P. K. SINHA, SANNIDHYA VARMA

Prevalence of Attention Deficit Hyperactivity Disorder in primary school children

JYOTHSNA AKAM VENKATA, ANUJA S. PANICKER

Post-stroke depression and lesion location –

A hospital based cross-sectional study

POOJA RAJASHEKARAN, KESHAVA PAI, RAVISH THUNGA, B. UNNIKRISHNAN

Efficacy of yoga for mental performance in university students

TIKHE SHAM GANPAT, HONGASANDRA RAMARAO NAGENDRA, V. SELVI

Prevalence and pattern of psychiatric morbidity and health related quality of life in patients with ischemic heart disease in a tertiary care hospital  
SHINY JOHN

Psychiatric morbidity and marital satisfaction among spouses of men with alcohol dependence

M. KISHOR, LAKSHMI V. PANDIT, R. RAGURAM

Psychiatric morbidity in prisoners

VINOD KUMAR, USHA DARIA

Chromosome 18p11.2 harbors susceptibility marker – D18S452, for bipolar affective disorder

MUTAHAR ANDRABI, ARSHAD HUSSAIN, FOUZIA RASHID, SHEIKH OZAIR NISSAR, IDREES AYOUB SHAH, YASIR HASAN RATHER, WASEEM HASSAN AHANGAR, NAZIR AHMAD DAR

## BRIEF RESEARCH COMMUNICATION

The course and outcome of alcohol use disorders in

men in Goa – A population-based follow-up study  
ABHIJIT NADKARNI, BHARGAV BHAT, SHAH EBRAHIM, VIKRAM PATEL

376

## CASE REPORTS

Body dysmorphic disorder – Borderline category between neurosis and psychosis

K. RAMAN

380

Dothiepin-induced transient hypomania and extrapyramidal syndrome

AMIR INAMDAR, ASHISH AGARWAL, PANKAJ KATARIA, R. P. BENIWAL

383

A case of Hallervorden-Spatz disease presenting as catatonic schizophrenia

YOGESH PAWAR, GURVINDER KALRA, SUSHMA SONAVANE, NILESH SHAH

386

Emetophobia – A fear of vomiting

ABHIJEET D. FAYE, SUSHIL GAWANDE, RAHUL TADKE, VIVEK C. KIRPEKAR, SUDHIR H. BHAVE

390

## CME

Forensic evaluations in psychiatry

R. K. CHADDA

393

## ART & PSYCHIATRY

Music and Mind

SRAVANTI SANIVARAPU

400

## VIEW POINT

The rise of super (?sub)-specialties courses in psychiatry – Is India ready for it!

N. MANJUNATHA, MURALI THYLOTH, T. S. SATHYANARAYANA RAO

401

## LETTERS TO EDITOR

Dissociative identity disorder – An uncommon psychiatric disorder reported

SIVA SHANKAR PRIYA, NAMBI SIVA

403

Studies on posttraumatic stress disorder – Challenges ahead

PALLAVI SINHA, AMIT GARG, OM PRAKASH

403

Prolonged, self-administration of ultra-high doses of quetiapine

AMAR BAVLE, CHITTARANJAN ANDRADE

404

Alcohol use among in-patients of a medical college hospital in Delhi

VINITHA C. THILAKAN, SANJEEV K. RASANIA

405

Job stress among emergency nursing staff – A preliminary study

GURVINDER PAL SINGH

407

## BOOK REVIEW

Em and the big Hoom – A book review

ALOK SARIN

409

## COMMENTARY

Have you forgotten?

JERRY PINTO

411



## The course and outcome of alcohol use disorders in men in Goa: A population- based follow-up study

Abhijit Nadkarni<sup>1,2</sup>, Bhargav Bhat<sup>2</sup>, Shah Ebrahim<sup>3,4</sup>, Vikram Patel<sup>1,2</sup>

<sup>1</sup>Department of Population Health, London School of Hygiene and Tropical Medicine, London, UK, <sup>2</sup>Sangath, Goa, <sup>3</sup>South Asia Network for Chronic Disease, Public Health Foundation of India, New Delhi, India, <sup>4</sup>Department of Non-communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

### ABSTRACT

**Background:** Research on the natural history and long-term outcomes of alcohol use disorders (AUD) is important to guide health policy. However, attrition is a major challenge in longitudinal studies and can affect validity of findings.

**Materials and Methods:** A 4-year follow-up study was conducted on a randomly selected subgroup of a sample of men in rural and urban communities in Goa to assess attrition, preliminary estimates of AUD outcomes, and chronic disease risk factors.

**Results:** Overall attrition rate was 16.7% with a higher attrition in AUD compared to abstainers or casual drinkers. Incidence of AUD at 4 years was 12.8% while the persistence rate of AUD was 53.9%. A trend for poorer health and health behaviors and higher blood pressure was observed in AUD.

**Conclusion:** The results of this study suggest a relatively low attrition rate despite the long follow-up period and support the feasibility of conducting a definitive cohort study with the entire sample.

**Key words:** Alcohol use disorders, attrition, health outcomes, incidence

### INTRODUCTION

The pattern of drinking in India is characterized by relatively high abstention rates and high rates of AUD among those who do drink.<sup>[1,2]</sup> This epidemiological pattern is associated with high rates of alcohol-attributable mortality and prevalence of AUDs, relative to the per capita alcohol consumption.<sup>[1]</sup> However, AUDs remain low on the Indian health policy agenda<sup>[3]</sup> and this could partly be due to the limited evidence on the impact of AUDs on health and socioeconomic outcomes.<sup>[2,4]</sup>

Evidence on the natural history and long-term outcomes of AUDs is of value to health policy makers. Such studies in developed countries have reported high rates of persistence

of AUD and adverse health and social outcomes.<sup>[5-12]</sup> However, one of the challenges of conducting such studies is attrition. Attrition represents a potential threat of bias, if those who drop out of the study are systematically different from those who remain in the study, which is plausible in the context of AUD. There is now a general consensus that loss of contact with more than 30% of the original sample over the course of a cohort study is unacceptable and will compromise the validity of conclusions drawn.<sup>[13]</sup>

The goal of this study was to estimate the attrition rates and reasons for attrition in a randomly selected subsample of a community cohort to inform the feasibility and methodology of a definitive evaluation of the course and outcomes of AUD in this population. We also planned

**Address for correspondence:** Dr. Abhijit Nadkarni, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK.  
E-mail: abhijit.nadkarni@lshtm.ac.uk

How to cite this article: Nadkarni A, Bhat B, Ebrahim S, Patel V. The course and outcome of alcohol use disorders in men in Goa: A population- based follow-up study. Indian J Psychiatry 2013;55:376-9.

#### Access this article online

##### Website:

[www.indianjpsychiatry.org](http://www.indianjpsychiatry.org)

##### DOI:

10.4103/0019-5545.120573

#### Quick Response Code



to determine the participant's current drinking status, co-occurrence of chronic disease risk factors, health outcomes, and acceptability of collection of blood samples in future studies.

## MATERIALS AND METHODS

### Setting

The study was conducted in the northern district of the state of Goa, India. The urban study areas included two beach areas popular among tourists and one urban residential area in Bardez taluka. The rural study areas include six contiguous villages in Sattari and Bicholim talukas.

### Sample

In 2006, the Alcohol Research Group (USA) and Sangath, Goa, conducted the SAAHAS study, which included a cross-sectional survey in the populations mentioned above.<sup>[14,15]</sup> Of the 1899 males that were screened for SAAHAS, 1859 (97.9%) consented or had not explicitly refused consent for follow-up contact. Alcohol use was assessed using the AUDIT.<sup>[16]</sup> It has been validated in India<sup>[17]</sup> and also specifically in Goa.<sup>[18]</sup> The WHO prescribed cut-off of 8 on AUDIT was used for detecting AUD. Participants who drank alcohol below this cut-off were coded as casual drinkers. Self-report of not having consumed any alcohol in the past year was coded as being abstainer. We selected a random sample of 50 men (25 each from rural and urban communities), from each of these subcategories, using random number tables, giving a total sample size of 150 men. As these data was collected in 2010 our outcomes were assessed roughly 4 years following baseline measurement.

### Measurements

A questionnaire measuring attrition, alcohol use, and common chronic disease risk factors was developed, and translated into the vernacular, i.e., Konkani, Hindi, and Marathi.

Chronic disease risk factors included questions related to physical activity, disability, mobility, smoking, and self-reported anxiety or depression. Blood pressure was measured using a digital sphygmomanometer which met the standards of the International Validation Protocol of the European Society of Hypertension (ESH).<sup>[19]</sup> Weight was measured using digital weighing scales and height was measured using a portable stadiometer.

If the respondent was unavailable after five follow-up attempts, had migrated to a different area which was unreachable by the team, had died or refused to participate, then he was considered as lost to follow-up and the reason for refusal or reason for loss to follow-up was recorded.

### Ethics

Ethical approval was obtained from Sangath's Institutional

Review Board. Written informed consent was obtained from each respondent and men with AUD were offered feedback and advice about help-seeking from Sangath or the public health system.

### Analyses

Categorical variables were summarized in the form of frequencies and proportions with 95% confidence intervals (CI). All outcomes were analyzed separately for the three exposure groups. Analyses were performed using STATA 10.0 for Windows.

## RESULTS

### Attrition rates

The overall attrition was 16.7% (95% CI 11.1-23.6). The reasons for attrition were as follows: address could not be traced 3.3%, migrated out of study area 6%, refused consent 5.3% and unavailable after five follow-up attempts 2%. The attrition in the urban centers was 28% (n = 21, 95% CI 18.2-39.6) and in the rural centers it was 5.3% (n = 4, 95% CI 2.2-14.9),  $P < 0.001$ . The attrition according to drinking status at baseline was 12% (95% CI 4.5-24.3) in abstainers, 16% (95% CI 7.2-29.1) in casual drinkers and 22% (95% CI 11.5-36) in AUD.

### Drinking outcomes at follow-up

Of the 86 abstainers/casual drinkers at baseline 12.8% (95% CI 6.6-21.7) were classified as AUD at follow-up with a higher incidence among casual drinkers (6.8%, 95% CI 1.4-18.7 in abstinent vs 19.1%, 95% CI 8.6-34.1, in casual drinkers). Of those who had AUD at baseline 53.9% (95% CI 38.3-71.4) were classified as persistent AUD at follow-up.

### Chronic disease risk factors

Table 1 compares the various chronic disease risk factors between abstainers, casual drinkers and hazardous drinkers. Compared to abstainers or casual drinkers, a higher proportion of those with AUD had self-reported history of hypertension, diabetes mellitus, chest pain or discomfort, moderate to severe general body pain, and shortness of breath. A slightly higher proportion of those with AUD were current smokers compared to casual drinkers or abstainers. A higher proportion of those with AUD (3.1%) were not physically active compared to abstainers (2.2%) and casual drinkers (2.1%). A higher proportion of AUD had problems with daily activities and mobility compared to the other two groups. Participants with AUD reported higher levels of moderate-to-severe anxiety or depression compared to abstainers and casual drinkers. The mean diastolic BP in AUD was higher than in casual drinkers and abstainers. The mean systolic BP in casual drinkers (130.1; 95% CI 126.5-133.9) was higher than in AUD (129.8; 95% CI 125.3-134.2) and abstainers (124.1; 95% CI 119.5-128.6). The mean BMI in casual drinkers was higher than in AUD and abstainers.

**Table 1: Chronic diseases and risk factors**

Variable	Abstainers <i>N</i> =46 (36.8%) <i>n</i> (%), 95% CI)	Casual drinkers <i>N</i> =47 (37.6%) <i>n</i> (%), 95% CI)	Hazardous drinkers <i>N</i> =32 (25.6%) <i>n</i> (%), 95% CI)
Hypertension	1 (2.2, 0.1-11.5)	6 (12.8, 4.8-25.7)	5 (15.6, 5.3-32.8)
Heart problem	3 (6.5, 1.4-17.9)	1 (2.1, 0.05-11.3)	1 (3.1, 0.07-16.2)
Diabetes mellitus	2 (4.4, 0.5-14.8)	4 (8.5, 2.4-20.4)	4 (12.5, 3.5-29.0)
Chest pain or discomfort	9 (19.6, 9.4-33.9)	7 (14.9, 6.2-28.3)	7 (21.9, 9.3-40.0)
Shortness of breath	9 (19.6, 9.4-33.9)	13 (27.7, 15.6-42.6)	11 (34.4, 29.8-74.3)
Current smoker	8 (17.4, 7.8-31.4)	11 (23.4, 12.3-38.0)	9 (28.1, 13.7-46.7)
Moderate to severe general body pain	19 (41.3, 27.0-56.8)	16 (34.0, 20.9-49.3)	19 (59.4, 40.6-76.3)
Problems with mobility and/or daily activities	7 (15.2, 6.3-28.9)	6 (12.8, 4.8-25.7)	11 (34.4, 18.6-53.2)
Moderate to severe anxiety or depression	16 (34.8, 21.4-50.2)	12 (25.5, 13.9-40.3)	22 (68.8, 50.0-83.9)
Mean diastolic BP	77.6 (74.7-80.5)	81.3 (77.6-85)	86.1 (15.2, 80.6-91.6)
Mean BMI	22.6 (21.4-23.8)	24.6 (23.4-25.8)	22.9 (21.4-24.4)

BMI – Body mass index; BP – Blood pressure

### Other outcomes

A total of 96% of the pilot study participants were willing to participate in a further follow-up in 2 years time and 91.2% expressed a willingness to undergo blood tests as a part of follow-up.

### DISCUSSION

The overall attrition rate in our sample was less than 20% at 4 years, the highest attrition being in AUD and lowest in abstainers; higher attrition was also observed in urban communities. Thus, we had a reasonably good retention rate at 4 years compared to previous longitudinal studies of drug and alcohol use which have reported attrition rates from 13% to 25% at 1 year.<sup>[20,21]</sup> A meta-analysis of longitudinal psychoactive substance prevention studies reported that on average 78.3% of participants were retained at 6 months, 73.4% after a year and 71.% after 2 years, suggesting that the largest attrition tends to occur during the first 6 months of a study.<sup>[22]</sup> Our attrition rate could be further reduced if we had resources to trace individuals who had migrated.

The incidence of AUD in our sample at 4 years was 12.8%. The only community-based Indian study that we identified reported an annual incidence rate of alcohol use disorders to be 4.2%.<sup>[23]</sup> In our sample, the abstinence rate in AUD was only 5.1% at 4 years. In the context of limited health services and large treatment gaps for AUDs, this is likely to approximate the rate of untreated abstinence. We do not have comparable studies in Indian populations but studies in developed countries in untreated samples report remission rates of 2-3%<sup>[24,25]</sup> at 1 year to 30% at 15 years.<sup>[26]</sup>

A higher proportion of those with AUD had adverse health and lifestyle outcomes compared to the other two groups. Although these findings are consistent with other studies<sup>[18,27-37]</sup> our small sample size restricts any inferences that we can draw from them. However, these findings do show a clear trend that AUD at baseline is associated with long-term adverse health outcomes; these findings coupled with those of low attrition strengthen the case for

conducting a definitive longitudinal study exploring the long-term course and outcome of AUD.

### ACKNOWLEDGMENT

This study was supported by the Wellcome Trust through the South Asia Network for Chronic Diseases of the Public Health Foundation of India and the LSHTM. Vikram Patel is supported by a Wellcome Trust Senior Research Fellowship. Abhijit Nadkarni is supported by a Wellcome Trust Research Training Fellowship.

### REFERENCES

1. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 2009;373:2223-33.
2. Prasad R. Alcohol use on the rise in India. *Lancet* 2009;373:17-18.
3. Kohn R, Saxena S, Levav I. The treatment gap in mental health care. *Bull World Health Organ* 2004;82:858-66.
4. Caetano R, Laranjeira R. A 'perfect storm' in developing countries: economic growth and the alcohol industry. *Addiction* 2006;101:149-52.
5. Gual A, Bravo F, Lligona A, Colom J. Treatment for alcohol dependence in Catalonia: Health outcomes and stability of drinking patterns over 20 years in 850 patients. *Alcohol Alcohol*. 2009;44:409-15.
6. Mann K, Schäfer DR, Längle G, Ackermann K, Croissant B. The long-term course of alcoholism, 5, 10 and 16 years after treatment. *Addiction* 2005;100:797-805.
7. McCabe RJ. Alcohol dependent individuals sixteen years on. *Alcohol Alcohol* 1986;21:85-91.
8. O'Connor A, Daly J. Alcoholics. A twenty year follow-up study. *Br J Psychiatry* 1985;146:645-7.
9. Hyman M. Alcoholics 15 Years Later. *Ann N Y Acad Sci* 1976;273:613-23.
10. Finney J, Moos R. The long-term course of treated alcoholism: I. Mortality, relapse and remission rates and comparisons with community controls. *J Stud Alcohol* 1991;52:44-54.
11. Gerdner A, Berglund M. Mortality of treated alcoholics after eight years in relation to short-term outcome. *Alcohol Alcohol* 1997;32:573-9.
12. Duckitt A, Brown D, Edwards G, Oppenheimer E, Sheehan M, Taylor C. Alcoholism and the nature of outcome. *Br J Addict* 1985;80:153-62.
13. McLellan AT, Woody GE, Metzger D. Evaluating the effectiveness of addiction treatments: reasonable expectations, appropriate comparisons. In: Egerton JA, Fox DM, Leshner A, editors. *Treating drug abusers effectively*. Massachusetts: Blackwell Publishers; 1997.
14. Nayak MB, Bond JC, Cherpitel C, Patel V, Greenfield TK. Detecting alcohol-related problems in developing countries: A comparison of 2 screening measures in India. *Alcohol Clin Exp Res* 2009;33:2057-66.
15. Nayak MB, Kerr W, Greenfield TK, Pillai A. Not all drinks are created equal: Implications for alcohol assessment in India. *Alcohol Alcohol* 2008;43:713-8.
16. Saunders JB, Aasland OG, Babor TF, Fuente JR, Grant M. Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction* 1993;88:791-804.



17. Babu RS, Sengupta SN, Sateesh Babu R. A study of problem drinkers in a general hospital. *Indian J Psychiatry* 1997;39:13-7.
18. Silva MC, Gaunekar G, Patel V, Kukalekar DS, Fernandes J. The prevalence and correlates of hazardous drinking in industrial workers: A study from Goa, India. *Alcohol Alcohol* 2003;38:79-83.
19. Belghazi J, Feghali RN, Moussalem T, Rejdych M, Asmar RG. Validation of four automatic devices for self-measurement of blood pressure according to the International Protocol of the European Society of Hypertension. *Vasc Health Risk Manag* 2007;3:389-400.
20. Aldridge J, Parker H, Measham F. Drug trying and drug use across adolescence: A longitudinal study of young people's drug taking in two regions of northern England. *Drugs prevention advisory service paper 1*. London: Home Office; 1999.
21. Botvin GJ, Baker E, Filazzola AD, Botvin EM. A cognitive-behavioral approach to substance abuse prevention: One-year follow-up. *Addict Behav* 1990;15:47-63.
22. Hansen WB, Tobler NS, Graham JW. Attrition in substance abuse prevention research: A meta-analysis of 85 longitudinally followed cohorts. *Eval Rev* 1990;14:677-85.
23. Mohan D, Chopra A, Sethi H. Incidence estimates of substance use disorders in a cohort from Delhi, India. *Indian J Med Res* 2002;115:128-35.
24. Gual A, Bravo F, Lligona A, Colom J. Treatment for alcohol dependence in Catalonia: Health outcomes and stability of drinking patterns over 20 years in 850 patients. *Alcohol Alcohol* 2009;44:409-15.
25. Vaillant GE. The natural history of alcoholism revisited. Cambridge, Massachusetts: Harvard University Press; 1995.
26. Öjesjö L. Long-term outcome in alcohol abuse and alcoholism among males in the Lundby general population, Sweden. *Br J Addict* 1981;76:391-400.
27. MacMahon S. Alcohol consumption and hypertension. *Hypertension* 1987;9:111-21.
28. Sisson JH. Alcohol and airways function in health and disease. *Alcohol* 2007;41:293-307.
29. Howard AA, Arnsten JH, Gourevitch MN. Effect of alcohol consumption on Diabetes Mellitus. *Ann Inter Med* 2004;140:211-9.
30. Paavola M, Vartiainen E, Haukkala A. Smoking, alcohol use, and physical activity: A 13-year longitudinal study ranging from adolescence into adulthood. *J Adolesc Health* 2004;35:238-44.
31. Battjes RJ. Smoking as an issue in alcohol and drug abuse treatment. *Addict Behav* 1988;13:225-30.
32. Di Franza JR, Guerreram P. Alcoholism and smoking. *J Stud Alcohol* 1990;51:130-5.
33. Hasin DS, Stinson FS, Ogburn E, Grant BF. Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: Results from the National Epidemiologic Survey on alcohol and related conditions. *Arch Gen Psychiatry* 2007;64:830-42.
34. Regier DA, Farmer ME, Rae DS, Locke BZ, Keith SJ, Judd LL, *et al.* Comorbidity of mental disorders with alcohol and other drug abuse: Results from the Epidemiologic Catchment Area (ECA) study. *JAMA* 1990;264:2511-8.
35. Upmark M, Hemmingsson T, Romelsjö A, Lundberg I, Allebeck P. Predictors of disability pension among young men: The role of alcohol and psychosocial factors. *Eur J Public Health* 1997;7:20-8.
36. D'Costa G, Nazareth I, Naik D, Vaidya R, Levy G, Patel V, *et al.* Harmful alcohol use in Goa, India, and its associations with violence: A study in primary care. *Alcohol Alcohol* 2007;42:131-7.
37. Gaunekar G, Patel V, Rane A. The impact and patterns of hazardous drinking amongst male industrial workers in Goa, India. *Soc Psychiatry Psychiatr Epidemiol* 2005;40:267-75.

**Source of Support:** Nil, **Conflict of Interest:** None declared